

SAT-LB-P-2-BFT(R)-09

USE OF ESSENTIAL OILS IN DAIRY PRODUCTS 4. ESSENTIAL OIL OF OREGANO (*ORIGANUM VULGARE* L.)

Assoc. Prof. Iliana Kostova, PhD

Department of Biotechnology and Food Technology, Razgrad Branch,
“Angel Kanchev” Univesity of Ruse
Phone: +359882836776
E-mail: ikostova@uni-ruse.bg

Prof. Stanka Damyanova, DcS

Department of Biotechnology and Food Technology, Razgrad Branch,
“Angel Kanchev” Univesity of Ruse
Phone: +359882669689
E-mail: sdamianova@uni-ruse.bg

Assoc. Prof. Nastya Ivanova, PhD

Department of Biotechnology and Food Technology, Razgrad Branch,
“Angel Kanchev” Univesity of Ruse
Phone: +359882669661
E-mail: nastiav2001@yahoo.com

Prof. Albena Stoyanova, DcS

Department of Essential oils,
University of Food Technologies, Plovdiv, Bulgaria
Tel.: +359894337990
E-mail: aastst@abv.bg

***Abstract:** There are dairy products developed with an enriched composition through the addition of linseeds, sesameseeds, oat flakes, honey, and essential oil of oregano (*Origanum vulgare* L.).*

The effect of the additives on the process of the acidifying, syneresis, and the development of lactic acid bacteria was researched. It was found that they are good for the lactic acid process. The prepared products have very good organoleptic properties and can be successfully used for the purposes of functional food.

***Keywords:** dairy products, titratable acidity, syneresis, lactic acid bacteria*

INTRODUCTION

Food is important to maintain and protect human health. A lot of traditional products (milk, fruits, vegetables, etc.) contain components with potential health benefits. New ones based on these food are being developed in a way increasing or joining the useful components because of their benefits or desirable physiological effects. Today the functional food based on milk takes up two-thirds of the total volume of such foods on the market as dairy foods with naturally balanced composition of the essential nutrients such as proteins, fats, carbohydrates, minerals, and enzymes. In the lactic acid products might be used: probiotics, dietary fibers (soluble and insoluble), vitamins, mineral elements, polyunsaturated fatty acids, essential oils, antioxidants, inulin, lactulose, etc. (Betored, E. et al. 2011, Cardarelli, H. et al. 2007, Kajiwara, S. et al. 2002, Panesar, P. S. 2011, Roberfroid, M. B. 2002).

In our previous work (Damyanova, St. et al. 2011) dairy products containing linseed and sesame seeds, oat flakes, and honey were prepared and examined. It was found that the products

have the properties of functional food and the additives have positive effect on the development of lactic acid process.

Interest in recent years is the use of essential oils in food products in order to increase their absorption by the human body, to enrich the composition, to improve the aroma, to prolong the durability, etc. (Barros, C. et al. 2012, Cleff, M. B. et al. 2010, Deliu, I. et al. 2017, Georgiev, E. & Stoyanova, A. 2006, Gutierrez, J. et al. 2009, Kačániová, M. et al. 2012, Thabet, H. et al. 2014).

The results from our previous studies (Kostova, I. et al. 2014) show that the essential oil of oregano has got antimicrobial activity but it does not inhibit the development of the lactic acid bacteria in dairy starter cultures.

The purpose of this work is to develop and examine dairy products with enriched composition by adding linseed, sesame seeds, oat flakes, oregano oil and honey.

EXPOSITION

Materials and methods

The studies were conducted in laboratory conditions with cow milk which was obtained from Razgrad region, northeastern Bulgaria.

The physicochemical parameters of raw milk (fat, solids-nonfat (SNF), density, added water, protein) were determined by the milk analyzer EKOMILK Company BULTEH 2000.

The microbiological parameters of the raw and pasteurized milk were tested by conventional methods (Slavchev, D. et al. 2003).

The fermented milk product with set coagulum (control) was prepared by a classical technology with the symbiotic starter culture of the strains *Lactobacillus delbrueckii subsp. bulgaricus* and *Streptococcus thermophilus* purchased by the company Selur Pharma Ltd.

The control sample was obtained by traditional technology for yogurt (Dimitrov, T. et al. 2008).

The samples studied (№ 1 to 3) were prepared by traditional technology by adding: oat flakes (6%), linseed (4%), sesame seed (2%) and honey (4%) (№ 1); essential oil (№ 2); the combination of the ingredients of the first and second examined samples (№ 3).

The additives used were market purchased; the essential oil of oregano was prepared in laboratory conditions (Kostova, I. et al. 2014). The amount of the oil is 0.8 mg/kg of final product, which was consistent with the known literal data and our previous studies (Georgiev, E. & Stoyanova, A. 2006, Kostova, I. et al. 2014).

The prepared dairy products were analyzed for chemical, microbiological and organoleptic characteristics:

- The lactic acid process dynamics was monitored by determination of titratable acidity (°T) (Dimitrov, T. et al. 2008, Slavchev, D. et al. 2003).

- Microbiological research - total number of viable lactic acid bacteria - *Lactobacillus delbrueckii subsp. bulgaricus* and *Streptococcus thermophilus* were carried out by growing on the synthetic medium M17 and MRS (Merck) methodology of IDF (IDF-Standard 122C: 1996, IDF-Standard 149A: 1997).

- The organoleptic assessment of lactic acid products was carried out according to BNS 15612-83 (BNS (Bulgarian National Standard) 15612-83).

RESULTS AND DISCUSSION

The milk used for the experimental work meets the requirements of raw cow milk (Commission Regulation (EC) No853/2004, Regulation № 4 the Ministry of Agriculture and Food 2008). Figure 1 presents the dynamics of the lactic acid process.




Fig. 1. Dynamics of lactic acid formation

The experiments conducted show increasement of the titratable acidity for all four samples as a result of lactic acid bacteria metabolism. More active lactic process was found in the samples with additives when they were combined with oregano oil. For these samples the tendency of increased acidity was maintained until the end of study period of 336 hours (14 days of the storage). The titratable acidity was higher than the control. The acid formation in samples with added essential oil was the same as for the control sample.




Fig. 2. Syneresis of the fermented milk products

Figure 2 presents the results for the values of the syneresis of the prepared dairy products. The amount of the separated whey for 5 hours was highest in the control sample (44.6 mL) and in the milk with oregano oil (42 mL). Significantly less whey was separated in sample № 1 (6.4 mL) and in sample №3 (7.6 mL), because of the ability of oat flakes and seeds to connect the water that leads to reduction of the aqueous phase of the milk.

The effect of the additives on development of lactic acid bacteria *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* in the tested products are shown in Figures 3 and 4.




Fig. 3. The growth dynamics of *Lactobacillus delbrueckii* subsp. *bulgaricus*

It was obvious from the data obtained that with the exception of the dairy product with oregano oil, in all other samples during the first 24 hours the cells of *Lactobacillus delbrueckii* subsp. *bulgaricus* grow actively. After 1st day and to 14th day, the number of the viable cells was kept relatively the same as in the dairy products with additives, in combination with the oregano oil (No 1 and 3) it was higher in comparison with the control. This is probably confirmed with the higher content of the nutrients needed for the development of the lactic bacteria. The results obtained correspond to the prescribed values for the titratable acidity (Fig. 1).

In the samples examined the number of the viable cells of *S. thermophilus* was increased to 48th hour. Then the growth speed was kept relatively unchanged till the end of the study period and the number of cells slightly decreases (Fig. 4). The lactic acid bacteria *S. thermophilus* develops fast growth speed in the samples with the additives and they reach the highest number in combination of all additives and the essential oil (sample №3).

The oregano oil inhibits to a certain extent the development of lactic acid bacteria *L. delbrueckii* subsp. *bulgaricus* and *S. thermophilus* that is related to its antimicrobial activity (Georgiev, E. & Stoyanova, A. 2006).




Fig. 4. The dynamics of cell growth of *Streptococcus thermophilus*

The data from the qualification of the prepared dairy products according to their organoleptic parameters are presented in Table 1. The products obtained were characterized as very good and balanced lactic acid taste, thick and smooth coagulum, homogeneous consistence and structure.

Table 1. Organoleptic indicators

Parameters	Control Sample	№ 1	№ 2	№ 3
Surface	smooth	smooth	smooth	smooth
Colour	white with cream-coloured shade	white with cream-coloured shade	white with cream-coloured shade	white with cream-coloured shade
Type of coagulum	thick, smooth	thick, smooth	thick, smooth	thick, smooth
Structure after cutting	smooth surface with slight serum separation	smooth surface without serum separation	smooth surface with slight serum separation	smooth surface without serum separation
Consistency after whipping the coagulum	homogeneous, like cream	homogeneous, like cream	homogeneous, like cream	homogeneous, like cream
Taste and flavor	tasty specific lactic acid	tasty specific lactic acid with a hint of the used nuts and honey	tasty specific lactic acid with a hint of oreganf	tasty specific lactic acid with a hint of the used nuts and honey, and oregano

CONCLUSION

Dairy products with added oat flakes, sesame seeds, linseeds, honey and essential oil of oregano (*Origanum vulgare* L.) has been prepared. The combination of these additives affects adversely on the speed of lactic acid process, syneresis, and their organoleptic parameters. The products acquired the specific for the fruits of oregano taste and aroma.

REFERENCES

- Barros, C., da Conceicao M., Gomes N., da Costa A., & da Souza, E. (2012). Combination of *Origanum vulgare* L. essential oil and lactic acid to inhibit *Staphylococcus aureus* in meat broth and meat model. *Brazilian Journal of Microbiology*, 1120 – 1127.
- Betored, E., Betored, N., Vidal, D., & Fito, P. (2011). Functional foods development: Trends and technologies. *Trends in Food Science & Technology*, 22, 498 – 508.
- BNS (Bulgarian National Standard) 15612-83 Dairy products. Organoleptic assessment.
- Cardarelli, H., Saad, S., Gibson, G., & Vulevich, J. (2007). Functional petit – suisse cheese: measure of the prebiotic effect. *Anaerobe*, 13, 200 – 207.
- Cleff, M. B., Meinerz, A. R., Xavier, M., Schuch, L. F., Meireles, M. C. A., Rodrigues, M. R. A., & Mello, J. R. B. (2010). *In vitro* activity of *Origanum vulgare* essential oil against *Candida* species. *Brazilian Journal of Microbiology*, 41, 116-123
- Commission Regulation (EC) No853/2004 (29.04.2004) laying down specific hygiene rules for on the hygiene of foodstuffs. *Off. Journal of European Union*, L139, 30.04.2004
- Damyanova, St., Todorova, S., Ivanova, N., Ganeva, E., & Stefanova, R. (2011). Development and testing of milk products with a functional purpose. *Kharkov National University Kiev NUHT*, 37, 47-51.
- Deliu, I., C. Nedelcu, M., & Teasă, C. M. (2017). Antimicrobial potential of garlic and oregano extracts and essential oils against different bacterial strains. *Current Trends in Natural Sciences*, 6 (12), 88-92
- Dimitrov, T., Mikhailova, G., Iliev, T., & Naydenova, N. (2008). Milk and milk products with research methods, Stara Zagora.

Georgiev, E., & Stoyanova, A. (2006). A guide for the specialist in aromatic industry, BNAEOPC, Plovdiv.

Gutierrez J., Barry-Ryan, C., Bourke, P. (2009). Antimicrobial activity of plant essential oils using food model media: efficacy, synergistic potential and interaction with food components. *Food Microbiology*, 26, 142-150.

IDF-Standard 122C: 1996 - Milk and milk products - Preparation of samples and dilutions for microbiological examination.

IDF-Standard 149A: 1997 - Dairy starter cultures of lactic acid bacteria - Standard of identity.

Kačániová, M., Vukovič, N., Hleba, L., Bobková, A., Pavelková, A., Rovná, K., & Arpášová, H. (2012). Antimicrobial and antiradicals activity of *Origanum vulgare* L. and *Thymus vulgaris* essential oils. *Journal of Microbiology, Biotechnology and Food Sciences*, 2 (1), 263-271

Kajiwara, S., Gandhi, H., & Ustunol, Z. (2002). Effect of honey on the growth of and acid production by human intestinal *Bifidobacterium spp.*: an *in vitro* comparison with commercial oligosaccharides and inulin. *J. Food Protect.*, 65, 214–218.

Kostova, I., Dimitrov, D., Ivanova, M., Vlaseva, R., Damyanova, S., Ivanova, N., Stoyanova, A. (2014). Studying the possibilities of using of essential oils in dairy products. *Oregano (Origanum vulgare L.) - Proceedings of the International Conference Modern Technologies, in the Food Industry – October 16-18, Chişinău (Republic of Moldova)*, 228-235.

Panesar, P. S. (2011). Fermented dairy products: Starter cultures and potential nutritional benefits. *Food and Nutrition Sciences*, 2, 47 – 51.

Regulation № 4 the Ministry of Agriculture and Food for the specific requirements for production, storage and transportation of raw cow milk and the requirements for trading and marketing of milk and milk products (Prom. SG. Br.23 of 29 February 2008). 2008.

Roberfroid, M. B. (2002). Global view on functional foods: European perspectives. *British Journal of Nutrition*, 88, Suppl. 2, S133 – S138.

Slavchev, D., Enikova, R., Makaveeva, M. 2003. Guide for physical, chemical and microbiological control of dairy products, Sofia.

Thabet, H., Nogaim, Q., Qasha, A., & Abdoalaziz, O. (2014). Evaluation of the effects of some plant derived essential oils on shelf life extension of Labneh. *Merit Research Journal of Food Science and Technology*, 21, 008 – 014.